



# Calcium6



## GUARANTEED ANALYSIS

Calcium (Ca) ..... 6.0%

DERIVED FROM: Calcium Carbonate

**CAUTION: KEEP OUT OF THE REACH OF CHILDREN, HARMFUL IF SWALLOWED**

9 lb/gal @ 68°F

## COMPATIBILITY

This product is compatible with most pesticides, however, it is not possible to test all pesticide mixes. A jar test is recommended before use.

## GENERAL USE RECOMMENDATIONS

**DILUTION:** Field and row crops: 3 to 20 gallons by air, 10 to 40 gallons by ground. Tree crops: concentrate 50 to 100 gallons, dilute 100 to 500 gallons.

## NET CONTENTS 2.5 GALLONS (9.46 L)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>



Guaranteed by KWS Distributing LLC.  
17887 SE Grand Island Rd.  
Dayton, Oregon 97114

## SUGGESTED USES: (rates per acre) Crops such as, but not limited to:

**Alfalfa, hay and other forage crops** :: 1 - 4 quarts after early spring growth and after cutting.

**Apples, cherries, peaches, almonds, walnuts, pears, stone, pome fruits, citrus and nuts** :: 1 - 4 quarts per acre at 2 to 4 week intervals.

**Barley, oats, wheat, rice, and other small grains** :: 1 - 4 quarts - Apply at tillering, early boot and flag leaf emergence.

**Beans, peas and other legumes** :: 1 - 3 quarts - Apply at early bud, repeat after bloom.

**Cabbage, broccoli, carrots, onions, tomatoes and other vegetables** :: 1 - 4 quarts - Apply when growth is 4 - 8" high. Repeat every 2 weeks as needed.

**Corn, field and sweet** :: 1 - 4 quarts - Apply when 10 - 20" high, repeat in 2 - 4 weeks as needed.

**Sugar Cane** :: 1 - 4 quarts per acre.

**Grapes** :: 1 - 4 quarts - Apply dormant, pre-bloom and post-bloom as needed.

**Hops** :: 1 - 4 quarts

**Lettuce, celery, spinach and other leafy vegetables** :: 1 - 4 quarts - Apply after transplanting or thinning, repeat as needed.

**Hazelnuts, pistachios and other deciduous trees** :: 1 - 4 quarts - Apply at bud break. Repeat every two to three weeks.

**Potatoes** :: 1 - 4 quarts - Apply at tuber initiation, repeat in 2 - 4 week intervals.

**Raspberries, blueberries, strawberries, blackberries** :: 1 - 3 quarts - Apply pre bloom repeat every 2-4 weeks.

**Grasses grown for seed** :: 1 - 4 quarts. Apply at tillering, early boot and flag leaf emergence.

**NOTICE** Manufacturer makes no warranties, expressed or implied. The buyer assumes all risks associated with the use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept a refund of the purchase price as full discharge of the manufacturers liability.

Rev. 070517

**CULTIVACEGROWTH.COM**  
A KWS DISTRIBUTING COMPANY