



GUARANTEED ANALYSIS

Copper (Cu) 10.0%

DERIVED FROM: Copper Acetate

CAUTION: KEEP OUT OF THE REACH OF CHILDREN, HARMFUL IF SWALLOWED

This product is intended to prevent or correct a copper deficiency.

9.6 lb/gal @ 68°F

COMPATIBILITY

This product is compatible with most pesticides, however, it is not possible to test all pesticide mixes. A jar test is recommended before use.

GENERAL USE RECOMMENDATIONS

DILUTION: Field and row crops: 10 to 20 gallons by air, 10 to 40 gallons by ground. Tree crops: concentrate 50 to 100 gallons, dilute 100 to 500 gallons.

NET CONTENTS 2.5 GALLONS (9.46 L)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>



Guaranteed by KWS Distributing LLC.
17887 SE Grand Island Rd.
Dayton, Oregon 97114

SUGGESTED USES: (rates per acre) Crops such as, but not limited to:

Alfalfa, hay and other forage crops :: 1 - 4 pints after early spring growth and after cutting.

Apples, cherries, peaches, almonds, walnuts, pears, stone, pome fruits, citrus and nuts :: 1 - 4 pints per acre at 2 to 4 week intervals.

Barley, oats, wheat, rice, and other small grains :: 1 - 4 pints - Apply at tillering, early boot and flag leaf emergence.

Beans, peas and other legumes :: 1 - 3 pints - Apply at early bud, repeat after bloom.

Cabbage, broccoli, carrots, onions, tomatoes and other vegetables :: 1 - 4 pints - Apply when growth is 4 - 8" high. Repeat every 2 weeks as needed.

Corn, field and sweet :: 1 - 4 pints - Apply when 10 - 20" high, repeat in 2 - 4 weeks as needed.

NOTICE Manufacturer makes no warranties, expressed or implied. The buyer assumes all risks associated with the use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept a refund of the purchase price as full discharge of the manufacturers liability.

Rev. 070517

Sugar Cane :: 1 - 2 pints per acre.

Grapes :: 1 - 4 pints - Apply dormant, pre-bloom and post-bloom as needed.

Hops :: 1 - 4 pints

Lettuce, celery, spinach and other leafy vegetables :: 1 - 4 pints - Apply after transplanting or thinning, repeat as needed.

Hazelnuts, pistachios and other deciduous trees :: 1 - 4 pints - Apply at bud break. Repeat every two to three weeks.

Potatoes :: 1 - 4 pints - Apply at tuber initiation, repeat in 2 - 4 week intervals.

Raspberries, blueberries, strawberries, blackberries :: 1 - 3 pints - Apply pre bloom repeat every 2-4 weeks.

Grasses grown for seed :: 1-4 pints. Apply at tillering, early boot and flag leaf emergence.

CULTIVACEGROWTH.COM
A KWS DISTRIBUTING COMPANY