



**GUARANTEED ANALYSIS**

Iron (Fe) ..... 5.0%

**DERIVED FROM:** Ferrous Acetate

**CAUTION: KEEP OUT OF THE REACH OF CHILDREN, HARMFUL IF SWALLOWED**

*This product is intended to prevent or correct an iron deficiency.*

10.0 lb/gal @ 68°F

**COMPATIBILITY**

This product is compatible with most pesticides, however, it is not possible to test all pesticide mixes. A jar test is recommended before use.

**GENERAL USE RECOMMENDATIONS**

**DILUTION:** Field and row crops: 3 to 20 gallons by air, 10 to 40 gallons by ground. Tree crops: concentrate 50 to 100 gallons, dilute 100 to 500 gallons.

**NET CONTENTS 2.5 GALLONS (9.46 L)**

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.org/metals.html>



Guaranteed by KWS Distributing LLC.  
17887 SE Grand Island Rd.  
Dayton, Oregon 97114



**PRECAUTIONARY STATEMENTS**

**Warning :** Causes skin irritation. Causes eye irritation. May cause respiratory irritation.

**Prevention :** Wear protective gloves.

Wear eye or face protection. Avoid breathing vapor.

**Response :** IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.

Call a POISON CENTER or physician if you feel unwell.

**Storage :** Store locked up.

**SUGGESTED USES:** (rates per acre) Crops such as, but not limited to:

**Alfalfa, hay and other forage crops ::** 1 – 8 pints after early spring growth, and after each cutting.

**Apples, cherries, peaches, almonds, walnuts, pears, stone, pome fruits, citrus and nuts ::** 1 – 8 pints per acre at 2 to 4 week intervals.

**Barley, oats, wheat, rice, and other small grains ::** 1 – 8 pints - Apply at tillering, early boot and flag leaf emergence.

**Beans, peas and other legumes ::** 1 – 6 pints - Apply at early bud, repeat after bloom.

**Cabbage, broccoli, carrots, onions, tomatoes and other vegetables ::** 1 – 8 pints - Apply when growth is 4 – 8" high. Repeat every 2 weeks as needed.

**Corn, field and sweet ::** 1 – 8 pints - Apply when 10 – 20" high, repeat in 2 – 4 weeks as needed.

**Sugar Cane:** 1—4 pints per acre.

**Grapes ::** 1 – 4 pints - Apply dormant, prebloom and post-bloom as needed.

**Hops ::** 1 – 4 pints

**Lettuce, celery, spinach and other leafy vegetables ::** 1 – 8 pints - Apply after transplanting or thinning, repeat as needed.

**Hazelnuts, pistachios, other deciduous trees ::** 1 – 8 pints - Apply at bud break. Repeat every two trees to three weeks.

**Potatoes ::** 1 – 8 pints - Apply at tuber initiation, repeat in 2 – 4 week intervals.

**Raspberries, blueberries, strawberries, blackberries ::** 1 – 3 pints - Apply pre bloom repeat every 2-4 weeks.

**Grasses grown for seed and turf ::** 1 – 8 pints. Apply at tillering, early boot and flag leaf emergence.

**NOTICE** Manufacturer makes no warranties, expressed or implied. The buyer assumes all risks associated with the use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept a refund of the purchase price as full discharge of the manufacturers liability.

Rev. 070517

**CULTIVACEGROWTH.COM**  
A KWS DISTRIBUTING COMPANY