



GUARANTEED ANALYSIS

TOTAL NITROGEN (N)	2.0%	Copper (Cu)	1.0%
2% Ammoniacal Nitrogen		Manganese(Mn)	1.0%
Available Phosphate(P ₂ O ₅)	20.0%	Zinc(Zn)	1.0%

DERIVED FROM: Ammonium Hydroxide, Phosphoric Acid, Copper Acetate, Zinc Acetate, Manganese Acetate.

KEEP OUT OF THE REACH OF CHILDREN, HARMFUL IF SWALLOWED

10.8 lb/gal @ 68°F

COMPATIBILITY

This product is compatible with most pesticides, however, it is not possible to test all pesticide mixes. A jar test is recommended before use.

GENERAL USE RECOMMENDATIONS

DILUTION: Field and row crops: 3 to 20 gallons by air, 10 to 40 gallons by ground. Tree crops: concentrate 50 to 100 gallons, dilute 100 to 500 gallons.

PRECAUTIONARY STATEMENTS

Danger : Causes severe skin burns and eye damage. Very toxic to aquatic life. Harmful to aquatic life with long lasting effects.

Prevention : Wear protective gloves. Wear eye or face protection. Wear protective clothing. Avoid release to the environment. Wash hands thoroughly after handling



CONTINUED ON NEXT PANEL

NET CONTENTS 2.5 GALLONS (9.46 L)

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.org/metals.html>



Guaranteed by KWS Distributing LLC.
17887 SE Grand Island Rd.
Dayton, Oregon 97114

SUGGESTED USES: (rates per acre) Crops such as, but not limited to:

Apples, cherries, peaches, almonds, walnuts, pears, stone, pome fruits, citrus and nuts :: 2 - 4 pints per acre at 2 to 4 week intervals starting at bud swell thru blossom drop.

Barley, oats, wheat, rice, and other small grains :: 2 - 4 pints - Apply at early boot, flag leaf emergence and pollination.

Beans, peas and other legumes :: 1 - 3 pints - Apply at early bud, repeat after bloom.

Cabbage, broccoli, carrots, onions, tomatoes and other vegetables grown for seed :: 1 - 4 pints - Apply 2 weeks before bloom. Make 2 more applications 2 weeks apart.

Corn, field and sweet :: 1 - 4 pints - Apply when 10 - 20" high, repeat in 2 - 4 weeks as needed.

Cont'd. from first panel: Response : Collect spillage. IF INHALED: Remove person to fresh air and keep comfortable for breathing. Immediately call a POISON CENTER or physician. IF SWALLOWED: Immediately call a POISON CENTER or physician. Rinse mouth. Do NOT induce vomiting. IF ON SKIN (or hair): Take off immediately all contaminated clothing. Rinse skin with water or shower. Wash contaminated clothing before reuse. Immediately call a POISON CENTER or physician. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or physician.

Storage : Store locked up.

Disposal : Dispose of contents and container in accordance with all local, regional, national and international regulations.

NOTICE Manufacturer makes no warranties, expressed or implied. The buyer assumes all risks associated with the use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept a refund of the purchase price as full discharge of the manufacturers liability.

Sugar Cane: 1—4 pints per acre.

Grapes :: 1 - 4 pints - Apply dormant, pre-bloom and post-bloom as needed.

Hops :: 1 - 4 pints

Hazelnuts, pistachios, other deciduous trees :: 2 - 4 pints - Apply at bud differentiation and again in 2 to 3 weeks.

Potatoes :: 2 - 4 pints - Apply at tuber initiation, repeat in 2 - 4 week intervals.

Raspberries, blueberries, strawberries, blackberries :: 1 - 3 pints - Apply pre bloom repeat every 2-4 weeks.

Grasses grown for seed :: 2 - 4 pints. Apply at early boot, flag leaf emergence and pollination.

Rev. 070517

CULTIVACEGROWTH.COM
A KWS DISTRIBUTING COMPANY