

FOLIAR APPLICATIONS FOR WINTER WHEAT

- 1) LATE JANUARY OR EARLY FEBRUARY: 1 QT/ACRE 3 ACES AND 2 - 4 OUNCES OF BORON. IF WHEAT ON WHEAT, ADD 1 QT MANGANESE ACETATE. IF RUST IS A PROBLEM, ADD 1 -2 PINTS OF COPPER ACETATE AND INCREASE THE BORON TO 4 – 8 OUNCES.
- 2) LATE FEBRUARY OR EARLY MARCH: 1 PINT/ACRE 3 ACES, 1 QT/ACRE COPPER ACETATE, 2 – 4 OUNCES OF BORON/ACRE, 1 PINT TO 1 QT./ACRE KELPGROW. IF RUST IS A PROBLEM, FOLLOW INFO IN #1.
- 3) APRIL: 1 PINT/ACRE 3 ACES, 1 – 2 PINTS/ACRE COPPER ACETATE, 1 – 2 QTS/ACRE K-ACE, 2 – 4 OUNCES BORON.
- 4) MAY: 1 – 2 PINTS/ACRE 2-6-0+Mo., 1 – 2 PINTS/ACRE COPPER ACETATE, 1 – 2 QTS K-ACE, 2 – 4 OUNCES BORON, ½ TO 1# CYTO-KELP.
- 5) JUNE: 1 – 2 PINTS/ACRE 2-6-0+Mo., 1 -2 GAL./ACRE SRN 28, 1 – 2 PINTS K-ACE, 2 – 4 OUNCES BORON.